

ANALYTIC PROPERTIES OF FORCE-FREE JETS IN THE KERR SPACETIME – I

ZHEN PAN,¹ CONG YU,^{2,3}

Submitted to ApJ

ABSTRACT

Blandford-Znajek (BZ) mechanism describes a process extracting rotation energy from a spinning black hole (BH) via magnetic field lines penetrating the event horizon of central BH. In this paper, we present a perturbation approach to study force-free jets launched by the BZ mechanism, and its two immediate applications: (1) we present a high-order split monopole perturbation solution to the BZ mechanism, which accurately pins down the energy extraction rate $\dot{E}$ and well describes the structure of BH magnetosphere for all range of BH spins ($0 \leq a \leq 1$); (2) the approach yields an exact constraint for the monopole field configuration in the Kerr spacetime, $I = \Omega(1 - A_\phi^2)$, where A_ϕ is the ϕ -component of the vector potential of electromagnetic field, Ω is the angular velocity of magnetic field lines and I is the poloidal electric current. The constraint is of particular importance to benchmark the accuracy of numerical simulations.

Subject headings: gravitation – magnetic fields – magnetohydrodynamics

1. INTRODUCTION

The Blandford-Znajek mechanism (Blandford & Znajek 1977) is believed to be an efficient way to extract rotation energy out of a spinning BH, which operates in BH systems on all mass scales, from stellar-mass BHs of gamma ray bursts to supermassive BHs of active galactic nuclei. In the past decade, general relativistic magnetohydrodynamic simulations have shown a physical picture of jets launched by the BZ mechanism (Komissarov 2001, 2004a,b, 2005; Komissarov & McKinney 2007; McKinney & Gammie 2004; McKinney 2005; Tchekhovskoy et al. 2008, 2010, 2011; Tchekhovskoy & McKinney 2012; Penna et al. 2013; McKinney et al. 2013), which shows that purely poloidal magnetic fields, $\mathbf{B}_P$, appear around a non-rotating BH, while a spinning BH induces poloidal electric field $\mathbf{E}_P$ and toroidal magnetic field $\mathbf{B}_T$, thus an outward Poynting flux $\mathbf{E}_P \times \mathbf{B}_T$ is generated, and rotation energy of the spinning BH is extracted in the form of Poynting flux.

Independent of time-dependent simulations, Contopoulos et al. (2013); Nathanail & Contopoulos (2014) numerically solved the general relativistic Grad-Shafranov (GS) equation which governs the structure of the stationary force-free magnetosphere around a Kerr BH. It is known that the GS equation is a second-order differential equation of A_ϕ depending on the magnetic field angular velocity Ω and the poloidal electric current I . The GS equation was shown to be an eigenvalue problem with eigenfunction A_ϕ and eigenvalues I and Ω to be determined. To solve the GS equation self-consistently, eigenvalues I and Ω were adjusted to make magnetic fields smoothly cross *two* light surfaces. In this way, they confirmed the results of previous simulations for asymptotically monopole and paraboloidal magnetic field. But for the asymptotically uniform mag-

netic field, there is only *one* light surface, which is not sufficient for determining the *two* eigenvalues I and Ω , so they argued that there are infinitely many solutions for the asymptotically uniform field (Contopoulos et al. 2013; Nathanail & Contopoulos 2014; Yang et al. 2015). While time-dependent simulations conducted by different groups seem to converge to similar solutions (e.g., Komissarov (2005); Komissarov & McKinney (2007); Alic et al. (2012)). But different simulations did not precisely match, especially in the range of extreme spins, and there is no way to tell which simulation is more robust.

To settle down all these uncertainties and controversies, independent analytic works are of great value (Tanabe & Nagataki 2008; Pan & Yu 2014, 2015). Though Contopoulos et al. (2013) criticized that the analytic perturbation approach did not appreciate the critical role of light surfaces, it is no coincidence that previous perturbation solutions are consistent with simulations, especially for non-extremal Kerr BHs (e.g. McKinney & Gammie (2004); Tchekhovskoy et al. (2010)). In fact, perturbation solutions obtained are analytical and smooth, which certainly satisfy the requirement of magnetic field lines smoothly crossing the light surfaces. So perturbation approach is consistent with the eigenvalue approach proposed by Contopoulos et al. (2013), and the only difference is that the eigenvalue approach implements the smoothly crossing requirement explicitly, while the perturbation approach implements the requirement implicitly.

In this paper, we propose a perturbation approach to investigate the BZ mechanism analytically, which in principle enables us to obtain perturbation solutions accurate to any order of the BH spin. Following this approach, we present a high-order monopole perturbation solution to the BZ mechanism, and derive an exact constraint relation for monopole magnetic field, which provides a criteria for testing the accuracy of numerical simulations. The paper is organized as follows. Basic equations governing stationary axisymmetric force-free magnetosphere in the Kerr spacetime are summarized in Section 2. In

¹ Department of Physics, University of California, One Shields Avenue, Davis, CA, 95616, USA, zhpan@ucdavis.edu

² Yunnan Observatories, Chinese Academy of Sciences, Kunming 650011, China, cyu@ynao.ac.cn

³ Key Laboratory for the Structure and Evolution of Celestial Objects, Chinese Academy of Sciences, Kunming 650011, China

Section 3, we present our perturbation method and apply it to the monopole magnetic field. We present results in Section 4 and discussion in Section 5. We conclude in Section 6.

2. BASIC EQUATIONS

Stationary axisymmetric force-free electromagnetic fields in the Kerr spacetime are determined by three functions A_ϕ , $\Omega(A_\phi)$, $I(A_\phi)$. Non-trivial components of Faraday tensor $F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu$ in the Kerr-Schild coordinate (McKinney & Gammie 2004; Kerr 1963) could be expressed as follows (Blandford & Znajek 1977; McKinney & Gammie 2004; Pan & Yu 2014)

$$F_{r\phi} = -F_{\phi r} = A_{\phi,r}, F_{\theta\phi} = -F_{\phi\theta} = A_{\phi,\theta}, \quad (1)$$

$$F_{tr} = -F_{rt} = \Omega A_{\phi,r}, F_{t\theta} = -F_{\theta t} = \Omega A_{\phi,\theta}, \quad (2)$$

$$F_{r\theta} = -F_{\theta r} = \sqrt{-g} B^\phi, \quad (3)$$

where

$$B^\phi = -\frac{I\Sigma + (2\Omega r - a)\sin\theta A_{\phi,\theta}}{\Delta\Sigma\sin^2\theta}, \quad (4)$$

and $\Sigma = r^2 + a^2\cos^2\theta$, $\Delta = r^2 - 2r + a^2$, $\sqrt{-g} = \Sigma\sin\theta$. The energy conservation equation reads

$$-\Omega[(\sqrt{-g}F^{tr})_{,r} + (\sqrt{-g}F^{t\theta})_{,\theta}] + F_{r\theta}I'(A_\phi) + [(\sqrt{-g}F^{\phi r})_{,r} + (\sqrt{-g}F^{\phi\theta})_{,\theta}] = 0, \quad (5)$$

which is also known as the Grad-Shafranov (GS) equation.

In the Schwarzschild spacetime, $I = \Omega = 0$, the GS equation is simplified as

$$\mathcal{L}A_\phi = 0, \quad (6)$$

where the operator

$$\mathcal{L} \equiv \frac{1}{\sin\theta} \frac{\partial}{\partial r} \left(1 - \frac{2}{r}\right) \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial}{\partial\theta} \frac{1}{\sin\theta} \frac{\partial}{\partial\theta}. \quad (7)$$

The Green's function $G(r, \theta; r_0, \theta_0)$ to the operator $\mathcal{L}$ defined by

$$\mathcal{L}G(r, \theta; r_0, \theta_0) = \delta(r - r_0)\delta(\theta - \theta_0), \quad (8)$$

is available (Blandford & Znajek 1977; Petterson 1974). Note that the solution satisfies the boundary conditions that $G(r, \theta; r_0, \theta_0)$ is finite at $r = 2$ and approaches to zero at infinity.

3. PERTURBATION METHOD

It is notoriously difficult to exactly solve the GS equation (5) due to its nonlinearity. We adopt the perturbation technique to attack this problem. Blandford & Znajek (1977) first put forward the monopole perturbation solution up to order of $O(a^2)$ and Pan & Yu (2015) extended the solution to $O(a^4)$. In this paper, we adopt a generalized perturbation approach, which, in principle, enables us to obtain perturbation solutions to any high order. We find that truncation at the eighth-order is good enough to precisely match current state-of-the-art numerical simulations. This method can be applied to other type of field configuration as well, though tedious calculations are inevitable. For simplicity, we focus on the split-monopole field throughout this paper.

For monopole magnetic field, the Schwarzschild metric solution to the GS equation (5) writes as

$$\Omega_0 = 0, \quad I_0 = 0, \quad A_\phi = A_0 = -\cos\theta. \quad (9)$$

For corresponding Kerr metric solution, we define $\omega = \Omega(A_\phi)|_{r \rightarrow \infty}$, $i = I(A_\phi)|_{r \rightarrow \infty}$, and expand them in series,

$$\begin{aligned} A_\phi &= A_0 + a^2 A_2 + a^4 A_4 + O(a^6), \\ \omega &= a\omega_1 + a^3\omega_3 + a^5\omega_5 + O(a^7), \\ i &= ai_1 + a^3i_3 + a^5i_5 + O(a^7). \end{aligned} \quad (10)$$

As done by Pan & Yu (2015), Ω and I could be expressed in terms of ω and i respectively. With the above notations, the GS equation (5) could be decomposed as a set of linear equations

$$\mathcal{L}A_n(r, \theta) = S_n(r, \theta; i_{n-1}, \omega_{n-1}) \quad (n = 2, 4, 6, \dots). \quad (11)$$

These equations could be solved one by one as follows.

Accurate to $O(a^2)$, GS equation (5) is written as

$$\mathcal{L}A_2(r, \theta) = S_2(r, \theta; i_1, \omega_1). \quad (12)$$

Before solving the differential equation, i_1 and ω_1 need to be specified by two constraints. The Znajek regularity condition (Znajek 1977) which requires B^ϕ to be finite on horizon gives

$$i_1 = \sin^2\theta \left(\frac{1}{4} - \omega_1\right), \quad (13)$$

and the convergence constraint that perturbation solution $A_2(r, \theta)$ should be convergent from horizon to infinity gives (see Pan & Yu (2015) for details)

$$i_1 = \omega_1 \sin^2\theta. \quad (14)$$

Consequently,

$$i_1 = \omega_1 \sin^2\theta, \quad \omega_1 = \frac{1}{8}. \quad (15)$$

With i_1 and ω_1 specified, the source function $S_2(r, \theta)$ is determined, so we can write the solution $A_2(r, \theta)$ as an integral of the Green's function

$$A_2(r, \theta) = \int_2^\infty dr_0 \int_0^\pi d\theta_0 S_2(r_0, \theta_0) G(r, \theta; r_0, \theta_0). \quad (16)$$

In this way, we work out all the three functions, A_ϕ , $I(A_\phi)$ and $\Omega(A_\phi)$ up to $O(a^2)$. Refer to Blandford & Znajek (1977) for explicit form of $A_2(r, \theta)$, where on inner/outer boundary

$$A_2(2, \theta) = R \sin^2\theta \cos\theta, \quad \lim_{r \rightarrow \infty} A_2(r, \theta) = 0, \quad (17)$$

with $R = (6\pi^2 - 49)/72$.

We could obtain solutions up to $O(a^4)$ in a similar manner (Pan & Yu 2015). The Znajek horizon condition and the convergence constraint give

$$i_3 = \omega_3 \sin^2\theta, \quad \omega_3 = \frac{1 - 4R}{64} \sin^2\theta + \frac{1}{32}, \quad (18)$$

and the integral of Green's function gives

$$\begin{aligned} A_4(2, \theta) &= p \sin^2\theta C_1^{(\frac{3}{2})}(\cos\theta) + q \sin^2\theta C_3^{(\frac{3}{2})}(\cos\theta), \\ \lim_{r \rightarrow \infty} A_4(r, \theta) &= 0. \end{aligned} \quad (19)$$

where

$$\begin{aligned} C_1^{(\frac{3}{2})}(\cos \theta) &= 3 \cos \theta, \\ C_3^{(\frac{3}{2})}(\cos \theta) &= -\frac{15}{2} \cos \theta + \frac{35}{2} \cos^3 \theta, \end{aligned} \quad (20)$$

and

$$\begin{aligned} p &= -\frac{9\sigma}{14} + \frac{13\zeta(3)}{3920} - \frac{\pi^2(19385 + 774\pi^2)}{181440} + \frac{17929399}{7620480} \\ &\simeq 1.7 \times 10^{-2}, \\ q &= -\frac{6\sigma}{7} + \frac{321\zeta(3)}{9800} - \frac{\pi^2(48955 + 918\pi^2)}{201600} + \frac{2012505017}{508032000} \\ &\simeq 9.0 \times 10^{-4}, \end{aligned} \quad (21)$$

with

$$\begin{aligned} \sigma &= \Re \int_2^\infty \frac{2\text{Li}_2\left(\frac{r}{2}\right) \log\left(\frac{r}{2}\right)}{\left(1 - \frac{2}{r}\right) r^2} dr = 1.3529... \\ \zeta(3) &= \frac{1}{2} \int_0^\infty \frac{t^2}{e^t - 1} dt = 1.2020... \end{aligned} \quad (22)$$

where $\Re$ designates the real part and $\text{Li}_2(x)$ is the second-order Jonquiere's function.

Take one step further, we could obtain quantities of $O(a^5)$

$$i_5 = \omega_5 \sin^2 \theta, \quad \omega_5 = \frac{1}{64} (1 - 0.037 \sin^2 \theta + 0.155 \sin^4 \theta), \quad (23)$$

using the Znajek horizon condition and the convergence constraint.

4. RESULTS

The energy extraction rate defined by $\dot{E} \equiv -2\pi \int_0^\pi \sqrt{-g} T_t^r d\theta = 2\pi \int I(A_\phi) \Omega(A_\phi) dA_\phi$, where T_ν^μ is energy-momentum tensor of electromagnetic field, is calculated accurate to $O(a^6)$ as

$$\begin{aligned} \dot{E} &= 2\pi \int i(A_0) \omega(A_0) dA_0 \\ &= \frac{\pi}{24} (a^2 + 0.587a^4 + 0.355a^6). \end{aligned} \quad (24)$$

Note that the 6th-order correction to the energy extraction rate only depends on $i_{1,3,5}$ and $\omega_{1,3,5}$ which are determined by $A_{0,2,4}$. See Fig. 1 for the energy extraction rate accurate to different orders. For comparison, corresponding simulation results from Tchekhovskoy et al. (2010) are also plotted. We find that the analytic energy extraction rate of $O(a^6)$ better matches simulation results compared to the original second-order BZ solution, but still underestimates energy extraction rate by $\sim 30\%$ for extreme spins, which indicates that even higher orders should be included. To improve the performance of analytic solutions, one way is to expand them to higher order of spin. We follow an alternative approach in this paper. We aim to find perturbation expansion series of faster convergence.

Simulations and analytic works agree that, $\omega = \Omega_H/2$ is a good approximation, where $\Omega_H = a/(2+2\sqrt{1-a^2})$ is angular velocity of a Kerr BH. We expect a faster solution convergence in terms of Ω_H instead of a . Accurate to




FIG. 1.— The energy extraction rate $\dot{E}$ accurate to different orders.




FIG. 2.— Distribution of angular velocity of magnetic field lines ω accurate to $O(\Omega_H^6)$.

$O(\Omega_H^6)$, we have

$$\omega = \frac{1}{2} \Omega_H + 0.43 \Omega_H^3 \sin^2 \theta + \Omega_H^5 (-5.75 \sin^2 \theta + 2.48 \sin^4 \theta), \quad (25)$$

which is shown in Fig. 2. We see $\omega \simeq \Omega_H/2$ for $a \lesssim 0.9$ and ω decrease abruptly for $a \gtrsim 0.9$. This turn-over was seen in many numerical simulations and is analytically confirmed here for the first time. In addition, take the extreme spin case $a = 0.999$ for example, we see the substructure that ω decreases quickly then increases slowly with increasing angle θ in the range of $0 \leq \theta \leq \pi/2$. The substructure was also confirmed by previous simulations (Tchekhovskoy et al. 2010).

With ω expressed in terms of Ω_H , the energy extraction rate is expressed in terms of Ω_H as

$$\dot{E} = \frac{2\pi}{3} (\Omega_H^2 + 1.38 \Omega_H^4 - 11.09 \Omega_H^6). \quad (26)$$

There is a smart way to extend $\dot{E}$ to even higher order making use of the relation

$$\left(\frac{d\dot{E}}{d\Omega_H} \right)_{a=1} = \left(\frac{d\dot{E}}{da} \frac{da}{d\Omega_H} \right)_{a=1} = 0, \quad (27)$$

where we have used the empirical constraint that $d\dot{E}/da$ is finite and the relation $(da/d\Omega_H)_{a=1} = 0$. As a result, eighth-order energy extraction rate $\dot{E}$ can be written as

$$\dot{E} = \frac{2\pi}{3}(\Omega_H^2 + 1.38\Omega_H^4 - 11.09\Omega_H^6 + 6.26\Omega_H^8). \quad (28)$$

Note that the eighth-order correction here is in principle different from the true correction that would be obtained in the usual way by solving the GS equation. The energy extraction rate $\dot{E}$ accurate to different orders are shown in Fig. 1. It turns out that our analytic solution up to $O(\Omega_H^8)$ perfectly reproduces the result of simulations for the whole range of spins. Besides, the commonly used lowest-order approximation $\dot{E} = (2\pi/3)\Omega_H^2$ overestimates the energy extraction rate by $\sim 30\%$ for the extreme spin, because, it does not account for the abrupt decrease of ω from $\Omega_H/2$ for $a \gtrsim 0.9$.

5. DISCUSSION

We find that $i_n = \omega_n \sin^2 \theta$ for $n \leq 5$ for split-monopole magnetic field configuration. Generally, we could prove the relation hold for all positive integer n . Take the $n = 8$ case for example,

$$\begin{aligned} S_8 = & \sin^2 \theta \omega_1 (\sin^2 \theta \omega_7)_{,\theta} + \sin^2 \theta \omega_3 (\sin^2 \theta \omega_5)_{,\theta} \\ & + \sin^2 \theta \omega_5 (\sin^2 \theta \omega_3)_{,\theta} + \sin^2 \theta \omega_7 (\sin^2 \theta \omega_1)_{,\theta} \\ & - (i_1 i_{7,\theta} + i_3 i_{5,\theta} + i_5 i_{3,\theta} + i_7 i_{1,\theta}) + O(1/r), \end{aligned} \quad (29)$$

where we have explicitly listed all source terms of $O(1)$. According to the convergence requirement, the summation of these $O(1)$ terms vanishes, so we have $i_7 = \omega_7 \sin^2 \theta$. Similarly, we find $i_n = \omega_n \sin^2 \theta$ for all positive n . As a result, we have $I = \Omega \sin^2 \theta|_{r \rightarrow \infty}$. Note that both I and Ω are functions of A_ϕ , so I/Ω is also a function A_ϕ . We denote it as $I/\Omega \equiv f(A_\phi)$. This function can be readily determined at infinity. It is known that $I/\Omega|_{r \rightarrow \infty} = \sin^2 \theta$ and $A_\phi|_{r \rightarrow \infty} = -\cos \theta$, so we have $f(-\cos \theta) = \sin^2 \theta$ or $f(x) = 1 - x^2$. Finally, we arrive at an exact constraint relation

$$I = \Omega(1 - A_\phi^2), \quad (30)$$

which holds at all radii. Recently, Penna (2015) shows that our constraint relation (30) is equivalent to the outgoing boundary condition at infinity.

According to this constraint relation, we show that the distortions of poloidal lines and variations of angular velocity of magnetic field lines are in fact results of force balance of magnetic field. We start with the explanation of $A_{\phi,\theta}$ variation with BH spin: BH rotation induces a toroidal component of magnetic field $\mathbf{B}_T$, whose hoop stress exerts a polar-directed force on poloidal field lines. In response, poloidal field lines move towards the polar direction (see Fig. 3), thus the fraction of magnetic flux increases in the polar region and decreases in the equatorial region. In other word, $|A_{\phi,\theta}|$ increases in the polar region and decreases in the equatorial region due to hoop stress of toroidal field generated by BH rotation. Combining the constraint relation (30) with the Znajek regularity condition, Ω on horizon $r = r_+$ can be expressed as

$$\frac{\Omega}{\Omega_H - \Omega} = \left(\frac{\sin \theta A_{\phi,\theta}}{1 - A_\phi^2} \right) \left(\frac{2r_+}{r_+^2 + a^2 \cos^2 \theta} \right), \quad (31)$$




FIG. 3.— Illustration of distortions of poloidal field lines induced by BH rotation, where solid lines are poloidal field lines in Kerr spacetime and dashed lines are corresponding field lines in Schwarzschild spacetime.

so Ω is determined by distortions of poloidal field lines and the size of BH horizon. For simplicity, we consider Ω in two regions: the equatorial region $\theta \simeq \pi/2$ and the polar region $\theta \simeq 0$. For the former,

$$\left(\frac{\Omega}{\Omega_H - \Omega} \simeq \frac{2A_{\phi,\theta}}{r_+} \right)_{\theta \simeq \pi/2}. \quad (32)$$

The right-hand side of the above equation is proportional to $\mathbf{B}_R/\mathbf{B}_T$, where the toroidal field $\mathbf{B}_T \sim 1/r$ and radial field $\mathbf{B}_R \sim A_{\phi,\theta}/r^2$. Decreasing $A_{\phi,\theta}$ will suppress $\mathbf{B}_R$. To keep the force balance, Ω must decrease accordingly to suppress hoop stress. Decreasing r_+ will increase more $\mathbf{B}_R$ than $\mathbf{B}_T$. To keep the force balance, Ω must increase accordingly. In real situation, both r_+ and $A_{\phi,\theta}$ decrease with increasing spin a , and Ω/Ω_H is determined by the competition between the two. For the polar region, we expect vanishing distortions of poloidal field lines, due to vanishing toroidal field, then we have

$$\Omega|_{\theta=0} = \Omega_H/2, \quad (33)$$

for any BH spin. Similar arguments also apply to other field configurations, e.g., magnetic field lines moving towards polar region and the turning over of Ω/Ω_H were indeed seen in simulations of paraboloidal magnetic field (Tchekhovskoy et al. 2010).

6. CONCLUSIONS

A stationary axisymmetric force-free jet launched by the BZ mechanism is completely determined by three function, A_ϕ , $I(A_\phi)$ and $\Omega(A_\phi)$, which are related by the GS equation (5). We propose an analytic approach to self-consistently determine the three functions. As a specific application, we present a high-order split monopole perturbation solution to the BZ mechanism. The solution accurately describes the structures of BH magnetosphere and the energy extraction rates of BZ mechanism for the whole range of BH spins. The analytic approach also yields an exact constraint relation, $I = \Omega(1 - A_\phi^2)$, for the split monopole magnetic field in the Kerr spacetime. The constraint relation

provides a useful tool for testing accuracy of numerical simulations. Based on this constraint relation, we find that the distortions of magnetic field lines and the variations of angular velocity Ω of magnetic field lines are results of force balance between the poloidal and toroidal field.

CY is grateful for the support by the National Natural

Science Foundation of China (Grant 11173057, 11373064, 11521303), Yunnan Natural Science Foundation (Grant 2012FB187, 2014HB048). Part of the computation was performed at the HPC Center, Yunnan Observatories, CAS, China. This work made extensive use of the NASA Astrophysics Data System and of the **astro-ph** preprint archive at arXiv.org.

REFERENCES

- Alic, D., Moesta, P., Rezzolla, L., Zanotti, O., & Jaramillo, J. L. 2012, *ApJ*, 754, 36
- Blandford, R. D., & Znajek, R. L. 1977, *MNRAS*, 179, 433
- Contopoulos, I., Kazanas, D., & Papadopoulos, D. B. 2013, *ApJ*, 765, 113
- Kerr, R. P. 1963, *Physical Review Letters*, 11, 237
- Komissarov, S. S. 2001, *MNRAS*, 326, L41
- . 2004a, *MNRAS*, 350, 427
- . 2004b, *MNRAS*, 350, 1431
- . 2005, *MNRAS*, 359, 801
- Komissarov, S. S., & McKinney, J. C. 2007, *MNRAS*, 377, L49
- McKinney, J. C. 2005, *ApJL*, 630, L5
- McKinney, J. C., & Gammie, C. F. 2004, *ApJ*, 611, 977
- McKinney, J. C., Tchekhovskoy, A., & Blandford, R. D. 2013, *Science*, 339, 49
- Nathanail, A., & Contopoulos, I. 2014, *ApJ*, 788, 186
- Pan, Z., & Yu, C. 2014, *Arxiv*, astro-ph/1406.4936
- . 2015, *Phys. Rev. D*, 91, 064067
- Penna, R. F. 2015, *ArXiv e-prints*, arXiv:1504.00360
- Penna, R. F., Narayan, R., & Sądowski, A. 2013, *MNRAS*, 436, 3741
- Petterson, J. A. 1974, *Phys. Rev. D*, 10, 3166
- Tanabe, K., & Nagataki, S. 2008, *Phys. Rev. D*, 78, 024004
- Tchekhovskoy, A., & McKinney, J. C. 2012, *MNRAS*, 423, L55
- Tchekhovskoy, A., McKinney, J. C., & Narayan, R. 2008, *MNRAS*, 388, 551
- Tchekhovskoy, A., Narayan, R., & McKinney, J. C. 2010, *ApJ*, 711, 50
- . 2011, *MNRAS*, 418, L79
- Yang, H., Zhang, F., & Lehner, L. 2015, *ArXiv e-prints*, arXiv:1503.06788
- Znajek, R. L. 1977, *MNRAS*, 179, 457